

# Cones of Hyperplane Arrangements and the Varchenko-Gel'fand Ring

Galen Dorpalen-Barry

School of Mathematics  
University of Minnesota

Kansas Combinatorics Seminar  
October 30, 2020 🐻



# Table of contents

1. Hyperplane Arrangements and Cones
2. The Varchenko-Gel'fand Ring
3. The Associated Graded Ring
4. A worked example of the Theorem
5. Supersolvable Arrangements

# Hyperplane Arrangements and Cones


# Hyperplane Arrangements

This talk is about (central) arrangements of hyperplanes  $\mathcal{A} = \{H_1, \dots, H_n\}$  in a real vector space  $V \cong \mathbb{R}^m$ .



# Hyperplane Arrangements

This talk is about (central) arrangements of hyperplanes  $\mathcal{A} = \{H_1, \dots, H_n\}$  in a real vector space  $V \cong \mathbb{R}^m$ .



Some notation:

1.  $\mathcal{C}(\mathcal{A})$  is the collection of chambers of  $\mathcal{A}$ .
2.  $\mathcal{L}(\mathcal{A})$  is the set of nonempty intersection subspaces  $X = H_{i_1} \cap H_{i_2} \cap \dots \cap H_{i_k}$ .
3. We will view  $\mathcal{L}(\mathcal{A})$  as a poset under (reverse) inclusion and define the (*signless*) *Whitney numbers of the first kind* for  $\mathcal{A}$  to be



$$c_k(\mathcal{A}) := \sum_{\substack{X \in \mathcal{L}(\mathcal{A}): \\ \text{codim}(X)=k}} |\mu(V, X)|.$$

Their generating function, the *Poincaré polynomial* of  $\mathcal{A}$ , is  $\text{Poin}(\mathcal{A}, t) := \sum_k c_k(\mathcal{A}) t^k$ .

# Hyperplane Arrangements

## Example

Here is an arrangement  $\mathcal{A} = \{H_1, H_2, H_3\} \subseteq \mathbb{R}^2$  (left) together with the Hasse diagram of its intersection poset  $\mathcal{L}(\mathcal{A})$  (right).



# Cones in an Arrangement

## Definition

A *cone*  $\mathcal{K}$  of an arrangement  $\mathcal{A}$  is an intersection of (open) half spaces defined by some of the hyperplanes of  $\mathcal{A}$ .


# Cones in an Arrangement

## Definition

A *cone*  $\mathcal{K}$  of an arrangement  $\mathcal{A}$  is an intersection of (open) half spaces defined by some of the hyperplanes of  $\mathcal{A}$ .

## Example

Let's consider a cone  $\mathcal{K}$  defined by  $H_4$  and  $H_5$  in the following three-dimensional arrangement of which I've drawn an affine slice.






# Cones in an Arrangement

## Definition

A *cone*  $\mathcal{K}$  of an arrangement  $\mathcal{A}$  is an intersection of (open) half spaces defined by some of the hyperplanes of  $\mathcal{A}$ .

## Example

Let's consider a cone  $\mathcal{K}$  defined by  $H_4$  and  $H_5$  in the following three-dimensional arrangement of which I've drawn an affine slice.



# Cones in an Arrangement

As with arrangements, a cone  $\mathcal{K}$  in an arrangement  $\mathcal{A}$  has chambers, intersections, and a Poincaré polynomial.

# Cones in an Arrangement

As with arrangements, a cone  $\mathcal{K}$  in an arrangement  $\mathcal{A}$  has chambers, intersections, and a Poincaré polynomial.

1. The *chambers of  $\mathcal{K}$*  are the chambers  $\mathcal{C}(\mathcal{K}) \subseteq \mathcal{C}(\mathcal{A})$  strictly contained in  $\mathcal{K}$ .

# Cones in an Arrangement

As with arrangements, a cone  $\mathcal{K}$  in an arrangement  $\mathcal{A}$  has chambers, intersections, and a Poincaré polynomial.

1. The *chambers of  $\mathcal{K}$*  are the chambers  $\mathcal{C}(\mathcal{K}) \subseteq \mathcal{C}(\mathcal{A})$  strictly contained in  $\mathcal{K}$ .
2. The nonempty intersections  $\mathcal{L}^{int}(\mathcal{K}) \subseteq \mathcal{L}(\mathcal{A})$  whose intersection with  $\mathcal{K}$  is nonempty are called *interior intersections of  $\mathcal{K}$* , i.e.  $X \in \mathcal{L}^{int}(\mathcal{K})$  if  $X \cap \mathcal{K} \neq \emptyset$ .

# Cones in an Arrangement

As with arrangements, a cone  $\mathcal{K}$  in an arrangement  $\mathcal{A}$  has chambers, intersections, and a Poincaré polynomial.

1. The *chambers of  $\mathcal{K}$*  are the chambers  $\mathcal{C}(\mathcal{K}) \subseteq \mathcal{C}(\mathcal{A})$  strictly contained in  $\mathcal{K}$ .
2. The nonempty intersections  $\mathcal{L}^{\text{int}}(\mathcal{K}) \subseteq \mathcal{L}(\mathcal{A})$  whose intersection with  $\mathcal{K}$  is nonempty are called *interior intersections of  $\mathcal{K}$* , i.e.  $X \in \mathcal{L}^{\text{int}}(\mathcal{K})$  if  $X \cap \mathcal{K} \neq \emptyset$ .
3. The (*signless*) *Whitney numbers of the first kind for  $\mathcal{K}$*  are



$$c_k(\mathcal{K}) := \sum_{\substack{X \in \mathcal{L}^{\text{int}}(\mathcal{K}): \\ \text{codim}(X)=k}} |\mu(\hat{0}, X)|.$$

Their generating function, the *Poincaré polynomial of  $\mathcal{K}$* , is  $\text{Poin}(\mathcal{K}, t) := \sum_k c_k(\mathcal{K}) t^k$ .

# Cones in an Arrangement





## Example



# Cones in an Arrangement

## Example



Thus  $\text{Poin}(\mathcal{K}, t) = 1 + 3t + t^2$ .

# Zaslavsky's Theorem for cones

## Theorem (Zaslavsky, '77)

For a cone  $\mathcal{K}$  of an arrangement  $\mathcal{A}$  with intersection poset  $\mathcal{L}^{int}(\mathcal{K})$ , we have

$$\#\mathcal{C}(\mathcal{K}) = \sum_{X \in \mathcal{L}^{int}(\mathcal{K})} |\mu(V, X)| = \sum_{k=0}^n c_k(\mathcal{K})$$

where  $\mu(V, X)$  denotes the Möbius function of  $\mathcal{L}^{int}(\mathcal{K})$  and  $\{c_k(\mathcal{K})\}$  are the Whitney numbers of the cone  $\mathcal{K}$ .

In other words  $\#\mathcal{C}(\mathcal{K}) = [\text{Poin}(\mathcal{K}, t)]_{t=1}$ .



# Zaslavsky's Theorem for cones

## Theorem (Zaslavsky, '77)

For a cone  $\mathcal{K}$  of an arrangement  $\mathcal{A}$  with intersection poset  $\mathcal{L}^{\text{int}}(\mathcal{K})$ , we have

$$\#\mathcal{C}(\mathcal{K}) = \sum_{X \in \mathcal{L}^{\text{int}}(\mathcal{K})} |\mu(V, X)| = \sum_{k=0}^n c_k(\mathcal{K})$$



where  $\mu(V, X)$  denotes the Möbius function of  $\mathcal{L}^{\text{int}}(\mathcal{K})$  and  $\{c_k(\mathcal{K})\}$  are the Whitney numbers of the cone  $\mathcal{K}$ .

In other words  $\#\mathcal{C}(\mathcal{K}) = [\text{Poin}(\mathcal{K}, t)]_{t=1}$ .

This result is well-known when we take  $\mathcal{K}$  to be the full arrangement.



# Zaslavsky's Theorem for cones

## Example





# Zaslavsky's Theorem for cones

## Example



# Zaslavsky's Theorem for cones



## Example



Zaslavsky says: there are  $1 + 3 + 1 = 5$  chambers in this cone.

# Zaslavsky's Theorem for cones

## Example



Zaslavsky says: there are  $1 + 3 + 1 = 5$  chambers in this cone.

**Goal:** Construct a ring from  $\mathcal{K}$  whose Hilbert Series is  $\text{Poin}(\mathcal{K}, t)$ .

# The Varchenko-Gel'fand Ring

# The Varchenko-Gel'fand Ring of a Cone

## Definition

The *Varchenko-Gel'fand ring* of a cone  $\mathcal{K}$  is the collection of maps  $VG(\mathcal{K}) = \{f : \mathcal{C}(\mathcal{K}) \rightarrow \mathbb{Z}\}$  under pointwise addition and multiplication.


# The Varchenko-Gel'fand Ring of a Cone

## Definition

The *Varchenko-Gel'fand ring* of a cone  $\mathcal{K}$  is the collection of maps  $VG(\mathcal{K}) = \{f : \mathcal{C}(\mathcal{K}) \rightarrow \mathbb{Z}\}$  under pointwise addition and multiplication.

For every cone  $\mathcal{K}$ ,  $VG(\mathcal{K})$  has a  $\mathbb{Z}$ -basis of indicator functions of chambers in  $\mathcal{C}(\mathcal{K})$ , as shown in the example.

## Example





# The Varchenko-Gel'fand Ring of a Cone

Pick an orientation of  $\mathcal{A}$ . It's easy to see that the Varchenko-Gel'fand ring  $VG(\mathcal{K})$  of a cone  $\mathcal{K}$  is generated by Heaviside functions

$$x_i(C) = \begin{cases} 1 & \text{if } v \in H_i^+ \cap \mathcal{K} \\ 0 & \text{else} \end{cases} \quad \text{for } C \in \mathcal{C}(\mathcal{K})$$

for each hyperplane  $H_i \in \mathcal{L}^{\text{int}}(\mathcal{K})$ .


# The Varchenko-Gel'fand Ring of a Cone

Pick an orientation of  $\mathcal{A}$ . It's easy to see that the Varchenko-Gel'fand ring  $VG(\mathcal{K})$  of a cone  $\mathcal{K}$  is generated by Heaviside functions

$$x_i(C) = \begin{cases} 1 & \text{if } v \in H_i^+ \cap \mathcal{K} \\ 0 & \text{else} \end{cases} \quad \text{for } C \in \mathcal{C}(\mathcal{K})$$


for each hyperplane  $H_i \in \mathcal{L}^{\text{int}}(\mathcal{K})$ .

## Example




# The Varchenko-Gel'fand Ring of a Cone

## Example



We can write the basis element corresponding to any chamber as a product of Heaviside functions for its walls.



$$= (1 - x_2)x_3x_4 = (1 - x_2)x_3$$

Define a map  $\varphi : \mathbb{Z}[e_1, \dots, e_n] \rightarrow VG(\mathcal{K})$  via  $e_i \mapsto x_i$ .

Define a map  $\varphi : \mathbb{Z}[e_1, \dots, e_n] \rightarrow VG(\mathcal{K})$  via  $e_i \mapsto x_i$ .

- ▶ By the previous observation, this map is **surjective**.
- ▶  $I_{\mathcal{K}} := \ker \varphi$  has a nice description.

# Signed Circuits

Recall,

# Signed Circuits

Recall,

- ▶ We can choose a set of normal vectors for the hyperplanes of  $\mathcal{A}$  so that  $v_i$  is normal to  $H_i$ .

# Signed Circuits

Recall,

- ▶ We can choose a set of normal vectors for the hyperplanes of  $\mathcal{A}$  so that  $v_i$  is normal to  $H_i$ .
- ▶ A *circuit* of a collection of vectors is a minimal dependent set, i.e. a set of linear dependent vectors such that if any single vector is removed, the set is independent. We will view circuits as sets of indices, so that  $C \subseteq \{1, 2, \dots, n\}$ .



# Signed Circuits

Recall,

- ▶ We can choose a set of normal vectors for the hyperplanes of  $\mathcal{A}$  so that  $v_i$  is normal to  $H_i$ .
- ▶ A *circuit* of a collection of vectors is a minimal dependent set, i.e. a set of linear dependent vectors such that if any single vector is removed, the set is independent. We will view circuits as sets of indices, so that  $C \subseteq \{1, 2, \dots, n\}$ .
- ▶ We'll keep track of *signed circuits* where we write down the explicit linear relations

$$\sum_{c \in C} \alpha_c v_c = 0 \quad \text{for } \alpha_c \in \mathbb{R}$$

and we sort the elements of  $C$  into  $C^+$  and  $C^-$ , depending on whether  $\alpha_c > 0$  or  $\alpha_c < 0$ .

# Presenting the Varchenko-Gel'fand Ring


## Theorem (D.-B., '20+)

Let  $\mathcal{K}$  be a cone of a central arrangement  $\mathcal{A} = \{H_1, \dots, H_n\}$ <sup>1</sup>. Then  $VG(\mathcal{K}) \cong \mathbb{Z}[e_1, \dots, e_n]/I_{\mathcal{K}}$  where  $I_{\mathcal{K}}$  is generated by

1. (Idempotent)  $e_i^2 - e_i$  for  $i \in [n]$ ,
2. (Unit)  $e_i - 1$  for  $i \in [n]$  such that  $H_i$  is a wall of  $\mathcal{K}$ ,
3. (Circuit)  $\prod_{i \in C^+} e_i \prod_{j \in C^-} (e_j - 1) - \prod_{i \in C^+} (e_i - 1) \prod_{j \in C^-} e_j$  for signed circuits  $C = C^+ \cup C^-$ ,

This was proved in 1987 by Varchenko and Gel'fand for  $\mathcal{K} = V$ .

---

<sup>1</sup>An affine arrangement is the cone of a higher-dimensional central arrangement. 

# Presenting the Varchenko-Gel'fand Ring

## Theorem (D.-B., '20+)


Let  $\mathcal{K}$  be a cone of a central arrangement  $\mathcal{A} = \{H_1, \dots, H_n\}$ <sup>1</sup>. Then  $VG(\mathcal{K}) \cong \mathbb{Z}[e_1, \dots, e_n]/I_{\mathcal{K}}$  where  $I_{\mathcal{K}}$  is generated by

1. (Idempotent)  $e_i^2 - e_i$  for  $i \in [n]$ ,
2. (Unit)  $e_i - 1$  for  $i \in [n]$  such that  $H_i$  is a wall of  $\mathcal{K}$ ,
3. (Circuit)  $\prod_{i \in C^+} e_i \prod_{j \in C^-} (e_j - 1) - \prod_{i \in C^+} (e_i - 1) \prod_{j \in C^-} e_j$  for signed circuits  $C = C^+ \cup C^-$ ,

This was proved in 1987 by Varchenko and Gel'fand for  $\mathcal{K} = V$ .

This theorem is actually a corollary to a stronger theorem...

---

<sup>1</sup>An affine arrangement is the cone of a higher-dimensional central arrangement. 

# Presenting the Varchenko-Gel'fand Ring

## Theorem (D.-B., '20+)

Let  $W = \{i \in [n] \mid H_i \text{ is a wall of } \mathcal{K}\}$ . For any graded monomial ordering on  $\mathbb{Z}[e_1, \dots, e_n]$ ,  $I_{\mathcal{K}}$  has Gröbner basis<sup>2</sup>:

1. (Idempotent)  $e_i^2 - e_i$  for  $i \in [n]$ ,
2. (Unit)  $e_i - 1$  for  $i \in [n]$  such that  $i \in W$
3. (Combination Circuit) Let  $C = C^+ \cup C^-$  be a signed circuit.
  - ▶ If  $W \cap C^\pm \neq \emptyset$  but  $W \cap C^\mp = \emptyset$ , then

$$\prod_{i \in C^+ \setminus W} e_i \prod_{j \in C^-} (e_j - 1) = \prod_{i \in C \setminus W} e_i - \pm \text{l.o.t.}$$

- ▶ If  $W \cap C = \emptyset$ , then

$$\prod_{i \in C^+} e_i \prod_{j \in C^-} (e_j - 1) - \prod_{i \in C^+} (e_i - 1) \prod_{j \in C^-} e_j = \sum_{j \in C} \pm \prod_{i \in C - \{j\}} e_i \pm \text{l.o.t.}$$

---

<sup>2</sup>The leading term of any polynomial in  $I_{\mathcal{K}}$  is divisible by the leading term of some polynomial in the Gröbner basis.

# No Broken Circuit Sets

Recall

# No Broken Circuit Sets

## Recall

- ▶ Let  $C$  be a circuit of  $\mathcal{A}$ . We can *break*  $C$  by removing the smallest index  $i$  contained in  $C$ . We call  $C - \{i\}$  the *broken circuit* corresponding to  $C$ .

# No Broken Circuit Sets

## Recall

- ▶ Let  $C$  be a circuit of  $\mathcal{A}$ . We can *break*  $C$  by removing the smallest index  $i$  contained in  $C$ . We call  $C - \{i\}$  the *broken circuit* corresponding to  $C$ .
- ▶ Let  $NBC(\mathcal{A})$  be the set of subsets of  $\{1, \dots, n\}$  containing no broken circuits.

# No Broken Circuit Sets

## Recall

- ▶ Let  $C$  be a circuit of  $\mathcal{A}$ . We can *break*  $C$  by removing the smallest index  $i$  contained in  $C$ . We call  $C - \{i\}$  the *broken circuit* corresponding to  $C$ .
- ▶ Let  $NBC(\mathcal{A})$  be the set of subsets of  $\{1, \dots, n\}$  containing no broken circuits.



# No Broken Circuit Sets

Recall

- ▶ Let  $C$  be a circuit of  $\mathcal{A}$ . We can *break*  $C$  by removing the smallest index  $i$  contained in  $C$ . We call  $C - \{i\}$  the *broken circuit* corresponding to  $C$ .
- ▶ Let  $NBC(\mathcal{A})$  be the set of subsets of  $\{1, \dots, n\}$  containing no broken circuits.

## Definition

A set  $N \in NBC(\mathcal{A})$  is a  $\mathcal{K}$ -NBC set if

$$\bigcap_{i \in N} H_i \in \mathcal{L}^{\text{int}}(\mathcal{K}).$$

Denote the set of  $\mathcal{K}$ -NBC sets by  $NBC(\mathcal{K})$ .

# A Basis for the Varchenko-Gel'fand Ring

## Theorem (D.-B., '20+)

Let  $\mathcal{K}$  be a cone of a central arrangement  $\mathcal{A}$ . Then  $VG(\mathcal{K})$  has

$$\left\{ \prod_{i \in N} e_i \mid N \in NBC(\mathcal{K}) \right\}$$

as a  $\mathbb{Z}$ -basis.

This was proved in 1987 by Varchenko and Gel'fand for  $\mathcal{K} = V$ .

# A Basis for the Varchenko-Gel'fand Ring

## Theorem (D.-B., '20+)


Let  $\mathcal{K}$  be a cone of a central arrangement  $\mathcal{A}$ . Then  $VG(\mathcal{K})$  has

$$\left\{ \prod_{i \in N} e_i \mid N \in NBC(\mathcal{K}) \right\}$$

as a  $\mathbb{Z}$ -basis.

This was proved in 1987 by Varchenko and Gel'fand for  $\mathcal{K} = V$ .

## Example



This is cool, but not quite what we wanted. Remember our goal:

**Goal:** Construct a ring from  $\mathcal{K}$  whose Hilbert Series is  $\text{Poin}(\mathcal{K}, t)$ .

# The Associated Graded Ring

# The Associated Graded Ring

- ▶ For  $d \geq 0$ , define  $F_d := \mathbb{Z} \cdot \{\text{monomials of degree} \leq d\} \subseteq VG(\mathcal{K})$ .

# The Associated Graded Ring

- ▶ For  $d \geq 0$ , define  $F_d := \mathbb{Z} \cdot \{\text{monomials of degree} \leq d\} \subseteq VG(\mathcal{K})$ .
- ▶ This yields a filtration  $\mathcal{F}$  of  $VG(\mathcal{K})$ :  $F_0 \subseteq F_1 \subseteq F_2 \subseteq \cdots$

# The Associated Graded Ring

- ▶ For  $d \geq 0$ , define  $F_d := \mathbb{Z} \cdot \{\text{monomials of degree } \leq d\} \subseteq VG(\mathcal{K})$ .
- ▶ This yields a filtration  $\mathcal{F}$  of  $VG(\mathcal{K})$ :  $F_0 \subseteq F_1 \subseteq F_2 \subseteq \dots$
- ▶ From this filtration, we define the *associated graded ring* of  $VG(\mathcal{K})$ :

$$\text{gr}_{\mathcal{F}}(VG(\mathcal{K})) := \bigoplus_{d \geq 0} F_d / F_{d-1}$$

where we set  $F_{-1} = 0$ .



# The Associated Graded Ring

- ▶ For  $d \geq 0$ , define  $F_d := \mathbb{Z} \cdot \{\text{monomials of degree } \leq d\} \subseteq VG(\mathcal{K})$ .
- ▶ This yields a filtration  $\mathcal{F}$  of  $VG(\mathcal{K})$ :  $F_0 \subseteq F_1 \subseteq F_2 \subseteq \dots$
- ▶ From this filtration, we define the *associated graded ring* of  $VG(\mathcal{K})$ :

$$\text{gr}_{\mathcal{F}}(VG(\mathcal{K})) := \bigoplus_{d \geq 0} F_d / F_{d-1}$$

where we set  $F_{-1} = 0$ .

- ▶ The Hilbert series (or Hilbert-Poincaré Series) of  $\text{gr}_{\mathcal{F}}(VG(\mathcal{K}))$  is the formal power series

$$\sum_{d \geq 0} \text{rk}_{\mathbb{Z}}(F_d / F_{d-1}) t^d$$

# The Hilbert Series of $\mathrm{gr}_{\mathcal{F}}(VG(\mathcal{K}))$

Theorem (D.-B., '20+)

*The Hilbert series of  $\mathrm{gr}_{\mathcal{F}}(VG(\mathcal{K}))$  is  $\mathrm{Poin}(\mathcal{K}, t)$ .*

This was proved in 1987 by Varchenko and Gel'fand for  $\mathcal{K} = V$ .



# The Hilbert Series of $\text{gr}_{\mathcal{F}}(\text{VG}(\mathcal{K}))$

Theorem (D.-B., '20+)

The Hilbert series of  $\text{gr}_{\mathcal{F}}(\text{VG}(\mathcal{K}))$  is  $\text{Poin}(\mathcal{K}, t)$ .

This was proved in 1987 by Varchenko and Gel'fand for  $\mathcal{K} = V$ .

Example




The theorem says that the Hilbert series of  $\text{gr}_{\mathcal{F}}(\text{VG}(\mathcal{K}))$  is  $1 + 3t + t^2$ .

## A worked example of the Theorem

# Example Computation I


Consider the following cone



The cone has 5 chambers, so  $VG(\mathcal{K}) \cong \mathbb{Z}^5$ . Earlier we computed its Whitney numbers, which are  $(1, 3, 1)$ .

# Example Computation I

Consider the following cone



The cone has 5 chambers, so  $VG(\mathcal{K}) \cong \mathbb{Z}^5$ . Earlier we computed its Whitney numbers, which are  $(1, 3, 1)$ .

## Example Computation II

Let's write down the Gröbner basis for  $I_{\mathcal{K}}$ . The Idempotent and Unit relations are

$$e_1^2 - e_1, e_2^2 - e_2, e_3^2 - e_3, e_4^2 - e_4, e_5^2 - e_5$$

and  $e_4 - 1, e_5 - 1$  respectively.

## Example Computation II

Let's write down the Gröbner basis for  $I_{\mathcal{K}}$ . The Idempotent and Unit relations are

$$e_1^2 - e_1, e_2^2 - e_2, e_3^2 - e_3, e_4^2 - e_4, e_5^2 - e_5$$

and  $e_4 - 1, e_5 - 1$  respectively. In order to write down the Combination Circuit relations, we need to do some work. The signed circuits are on the left and the relation is on the right:

$$\begin{aligned}\{2, 5\} \cup \{1\} &\rightarrow e_2(e_1 - 1) = e_1 e_2 - e_2 \\ \{1, 3\} \cup \{2, 4\} &\rightarrow e_1 e_3(e_2 - 1) = e_1 e_2 e_3 - e_1 e_3 \\ \{3, 4, 5\} \cup \{1\} &\rightarrow (e_1 - 1)e_3 = e_1 e_3 - e_3 \\ \{2, 4\} \cup \{3, 5\} &\rightarrow 0\end{aligned}$$



## Example Computation III

From this we can write down the NBC-basis of  $VG(\mathcal{K})$  itself. The circuits are on the left and the broken circuits are on the right:

$$125 \rightarrow 25$$

$$1234 \rightarrow 234$$

$$1345 \rightarrow 345$$

$$2345 \rightarrow 345$$

The no broken circuit sets associated to  $\mathcal{A}$  are:

$\emptyset,$

1, 2, 3, 4, 5,

12, 13, 14, 15, 23, 24, 34, 35, 45,

123, 124, 134, 135, 145

## Example Computation III

From this we can write down the NBC-basis of  $VG(\mathcal{K})$  itself. The circuits are on the left and the broken circuits are on the right:

$$125 \rightarrow 25$$

$$1234 \rightarrow 234$$

$$1345 \rightarrow 345$$

$$2345 \rightarrow 345$$

The  $\mathcal{K}$ -NBC sets are:

$\emptyset,$


**1, 2, 3, 4, 5,**

12, 13, 14, 15, **23**, 24, 34, 35, 45,

123, 124, 134, 135, 145

# Example Computation IV

The NBC-basis for  $VG(\mathcal{K})$  is



So the associated graded ring is

$$\mathrm{gr}_{\mathcal{F}}(VG(\mathcal{K})) \cong \mathbb{Z} \cdot \{1\} \oplus \mathbb{Z} \cdot \{x_1, x_2, x_3\} \oplus \mathbb{Z} \cdot \{x_2x_3\}$$

and has Hilbert series  $1 + 3t + t^2$ .

# Supersolvable Arrangements


# What is a supersolvable arrangement?


## Definition

An arrangement is *supersolvable* if there is a maximal chain  $\Delta$  of the intersection lattice  $\mathcal{L}(\mathcal{A})$  such that for every chain  $K$ , the sublattice generated by  $\Delta$  and  $K$  is *distributive*<sup>3</sup>.

## Example

The  $(n - 1)$ st braid arrangement is supersolvable and consists of hyperplanes  $H_{ij} = \{\mathbf{x} \in \mathbb{R}^d \mid x_i = x_j\}$  for  $i, j \in [n]$ . A linearly equivalent picture of the  $(3 - 1)$ st braid arrangement is below (left) together with its intersection poset  $\mathcal{L}(\mathcal{A})$  (right).



<sup>3</sup>A lattice  $L$  is distributive if for all  $x, y \in L$ , we have  $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ . 



# What is a supersolvable arrangement?

## Theorem (Björner-Ziegler, '91)


When we order the broken circuits of a supersolvable arrangement by inclusion, the minimal broken circuits have cardinality exactly 2.

## Example

The  $(3 - 1)$ st braid arrangement.



There is one circuit consisting of all three hyperplanes  $\{1, 2, 3\}$ .  
The broken circuit is  $\{2, 3\}$ .


 The  $(n - 1)$ st braid arrangement is the *complete graph arrangement*.

**Upshot:** We can write down the circuits of the braid arrangement from the *circuits* of the complete graph.

# What does being supersolvable have to do with the Varchenko-Gel'fand ring?


## Definition

The *Varchenko-Gel'fand ring* of a cone  $\mathcal{K}$  over a field  $\mathbb{F}$  is the collection of maps  $VG_{\mathbb{F}}(\mathcal{K}) = \{f : \mathcal{C}(\mathcal{K}) \rightarrow \mathbb{F}\}$  under pointwise addition and multiplication.

 Our previous theorems still hold for  $VG_{\mathbb{F}}(\mathcal{K})$   
(in fact they are easier because we're now over a field!)

## Theorem (D.-B. '20+)

*If  $\mathcal{A}$  is a supersolvable arrangement, then for every cone  $\mathcal{K}$ , the associated graded ring  $\text{gr}(VG_{\mathbb{F}}(\mathcal{K}))$  is Koszul.*

 This theorem fits into a larger context.

# Fitting this into a Larger Context: the Orlik-Solomon Algebra

The Orlik-Solomon algebra is a noncommutative analogue of the Varchenko-Gel'fand ring.

## Theorem (D.-B. '20+)

*If  $\mathcal{A}$  is a supersolvable arrangement, then for every cone  $\mathcal{K}$ , the associated graded ring  $\mathrm{gr}(VG_{\mathbb{F}}(\mathcal{K}))$  is Koszul.*

## Theorem (Peeva '02)

*If  $\mathcal{A}$  is a supersolvable arrangement, then the Orlik-Solomon algebra of  $\mathcal{A}$  is supersolvable.*



Thank you!

# Selected References I



William W. Adams and Philippe Loustau.

*An introduction to Gröbner bases*, volume 3 of *Graduate Studies in Mathematics*.

American Mathematical Society, Providence, RI, 1994.



Anders Björner, Michel Las Vergnas, Bernd Sturmfels, Neil White, and Günter M. Ziegler.

*Oriented matroids*, volume 46 of *Encyclopedia of Mathematics and its Applications*.

Cambridge University Press, Cambridge, second edition, 1999.



A. N. Varchenko and I. M. Gel'fand.

Heaviside functions of a configuration of hyperplanes.

*Funktsional. Anal. i Prilozhen.*, 21(4):1–18, 96, 1987.



Thomas Zaslavsky.

A combinatorial analysis of topological dissections.

*Advances in Math.*, 25(3):267–285, 1977.